

Renal injury in extreme obesity: the important role of aldosterone

Kidney International (2008) **74**, 1216; doi:10.1038/ki.2008.429

To the Editor: We read with interest the recent article by Serra *et al.*¹ detailing a variety of glomerular lesions (increased mesangial matrix, mesangial cell proliferation, podocyte hypertrophy, and glomerulomegaly) in extremely obese patients without clinical signs of renal dysfunction. Despite using a broad array of clinical and biochemical variables in their regression models, only body mass index was found to be a significant predictor of glomerular lesions in multivariate analysis.

We believe that serum aldosterone, if included in the analysis, would have likely emerged as another important predictor of obesity-related renal damage. Obesity is frequently associated with elevated levels of aldosterone.² In the obese state, oxidized fatty acids likely stimulate aldosteronogenesis;³ *in vitro*, human adipocytes secrete potent mineralocorticoid-releasing factors.⁴ Aldosterone is important in salt and water balance by acting on epithelial mineralocorticoid receptors, but its effects through mineralocorticoid receptors in nonepithelial tissues may be more important in the pathogenesis of chronic kidney disease. In animal models, unopposed aldosterone in the presence of high salt intake causes thrombotic and proliferative lesions in the glomeruli and renal vessels.^{5–7} These pathologic lesions occur independent of blood pressure, reflecting a direct, nonepithelial, profibrotic effect of aldosterone on the kidney.

The increased glomerulosclerosis in these hyperaldosterone models typically result in severe proteinuria, a distinctly different phenotype from Serra *et al.*'s extremely obese subjects whose clinical renal function was normal. The authors, however, argue that the early lesions found in this study are potential harbingers of future, overt kidney disease. If so, then the glomerular lesions described in this study will likely progress to the more sclerotic lesions typified in the seminal animal studies discussed above.

The role of aldosterone in obesity-related kidney injury becomes more important as the discussion shifts toward potential treatment options. Serra *et al.* admit that their study does not address whether bariatric surgery could reverse the renal lesions seen on biopsy, and the authors cite a study in which angiotensin-converting enzyme inhibitors and statins ameliorate podocyte damage in obese rats.⁸ In dogs fed a high-fat diet, the use of an aldosterone antagonist (compared to untreated controls) markedly attenuated obesity-induced glomerular hyperfiltration, sodium retention, and hypertension.⁹ Because it will not be possible to offer all extremely obese patients bariatric surgery, it may be beneficial to check these patients' aldosterone levels and, if elevated, prescribe mineralocorticoid receptor blockers and salt restriction for renal protection. Alternatively, empiric treatment with

mineralocorticoid receptor blockers and a low-salt diet may be undertaken even without aldosterone measurement, recognizing the possibility that mineralocorticoid receptor activation may also occur by elevated cortisol concentrations in obesity.¹⁰

1. Serra A, Romero R, Lopez D *et al.* Renal injury in the extremely obese patients with normal renal function. *Kidney Int* 2008; **73**: 947–955.
2. Goodfriend TL, Egan BM, Kelley DE. Aldosterone in obesity. *Endocr Res* 1998; **24**: 789–796.
3. Goodfriend TL, Ball DL, Egan BM *et al.* Epoxy-keto derivative of linoleic acid stimulates aldosterone secretion. *Hypertension* 2004; **43**: 358–363.
4. Ehrhart-Bornstein M, Lamounier-Zepter V, Schraven A *et al.* Human adipocytes secrete mineralocorticoid-releasing factors. *Proc Natl Acad Sci USA* 2003; **100**: 14211–14216.
5. Greene EL, Kren S, Hostetter TH. Role of aldosterone in the remnant kidney model in the rat. *J Clin Invest* 1996; **98**: 1063–1068.
6. Rocha R, Chander PN, Zuckerman A *et al.* Role of aldosterone in renal vascular injury in stroke-prone hypertensive rats. *Hypertension* 1999; **33**(1, Part 2): 232–237.
7. Rocha R, Stier Jr CT, Kifor I *et al.* Aldosterone: a mediator of myocardial necrosis and renal arteriopathy. *Endocrinology* 2000; **141**: 3871–3878.
8. Blanco S, Vaquero M, Gomez-Guerrero C *et al.* Potential role of angiotensin-converting enzyme inhibitors and statins on early podocyte damage in a model of type 2 diabetes mellitus, obesity, and mild hypertension. *Am J Hypertens* 2005; **18**(4, Part 1): 557–565.
9. de Paula RB, da Silva AA, Hall JE. Aldosterone antagonism attenuates obesity-induced hypertension and glomerular hyperfiltration. *Hypertension* 2004; **43**: 41–47.
10. Frey FJ, Odermatt A, Frey BM. Glucocorticoid-mediated mineralocorticoid receptor activation and hypertension. *Curr Opin Nephrol Hypertens* 2004; **13**: 451–458.

Andrew S. Bomback¹ and Philip J. Klemmer¹

¹*Division of Nephrology and Hypertension, Department of Medicine, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA*

Correspondence: Andrew S. Bomback, University of North Carolina Kidney Center, 7024 Burnett-Womack Building, Campus Box 7155, Chapel Hill, North Carolina 27599-7155, USA. E-mail: abomback@unch.unc.edu

Response to 'renal injury in extreme obesity: the important role of aldosterone'

Kidney International (2008) **74**, 1216–1217; doi:10.1038/ki.2008.438

We appreciate the comments of Dr Bomback and Dr Klemmer on the possible relationship between aldosterone and obesity-related renal lesions.¹

Obesity is associated with elevated levels of several hormones including aldosterone, leptin, adiponectin, cortisol, renin, and angiotensin.^{2,3} Some of these hormones, such as leptin, have been associated with the production of renal lesions in animal models.⁴